



Unwin, S. L., Saunders, R., Blackwell, E. J., & Rooney, N. J. (2019). A double-blind, placebo-controlled trial investigating the value of Pet Remedy in ameliorating fear of handling in companion rabbits. *Journal of Veterinary Behavior: Clinical Applications and Research*.  
<https://doi.org/10.1016/j.jveb.2019.10.001>

Peer reviewed version

License (if available):  
CC BY-NC-ND

Link to published version (if available):  
[10.1016/j.jveb.2019.10.001](https://doi.org/10.1016/j.jveb.2019.10.001)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://www.sciencedirect.com/science/article/pii/S1558787818301898#!>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

1     **A double-blind, placebo-controlled trial investigating the**  
2     **value of Pet Remedy in ameliorating fear of handling of**  
3     **companion rabbits**

4                     Authors: Unwin SL, Saunders R, Blackwell EJ, Rooney NJ.

7                     Short Title: Investigating the value of Pet Remedy use in companion rabbits

9                     Corresponding author: [Nicola.Rooney@bristol.ac.uk](mailto:Nicola.Rooney@bristol.ac.uk)

10                    Animal Welfare and Behavior Group, Bristol Veterinary School, University of Bristol, Langford, BS40

11                                     5DU

## Abstract

The domestic rabbit is a popular companion animal in the UK with an estimated population of 0.9 million. Research has highlighted a large number of welfare issues related to the way rabbits are commonly kept, affecting a majority of the population. One major welfare issue is the large proportion of pet rabbits that are fearful when handled. Pet Remedy™ (Unex Designs) is an herbal product containing valerian, marketed as a natural calming aid for reducing stress in all companion animal species. Its efficacy for domestic rabbits is previously untested. We describe a randomized, double-blind, placebo-controlled trial into the effectiveness of Pet Remedy when sprayed on an object in the environment and the clothing of a novel person handling the rabbit.

A rehoming population of 50 rabbits each underwent a baseline test followed by both a placebo and a Pet Remedy treatment on three consecutive days (following cross-over design). On each day, a novel arena test was conducted, and the rabbits' behavioral responses to the experimenter measured when in its home enclosure and when being handled. Heart and respiratory rates were measured during handling. Repeated measures ANOVAs tested within-subjects differences between Pet Remedy and placebo trials, while taking into account rabbits' individual baselines.

Treatment with Pet Remedy was associated with a significant decrease in heart rate during handling ( $F_{(1,42)}=4.41$ ,  $P=0.042$ ) and a significant increase in the number of positive behaviors observed in a novel arena test ( $F_{(1,47)}=4.52$ ,  $P=0.039$ ). Rabbits took marginally longer to be picked up at the start of a Pet Remedy trial ( $F_{(1,47)}=4.08$ ,  $P=0.049$ ). Other variables which may have been predicted to change were unaffected. Overall the amount of rearing in the novel arena increased with day ( $F_{(1,45)}=6.91$ ,  $P=0.012$ ), showing an increase in exploratory behavior with habituation. Significant individual variation occurred throughout, with almost all variables being markedly affected by an

individual's baseline data, and heart rates were universally high suggesting that handling is generally aversive to many rabbits.

The results of this study suggest that Pet Remedy may have some potential value for rabbits during periods of acute stress, slowing heart rate and allowing the performance of more positive, relaxed behaviors. It may thus be useful during veterinary visits and during initial handling. However, given the high levels of physiological and behavioral stress exhibited, effects were small and hence we suggest optimal handling and appropriate habituation, desensitisation and counter-conditioning protocols should also be simultaneously implemented. The efficacy of pet remedy for long-term use remains untested.

## Introduction

The domestic rabbit (*Oryctolagus cuniculus*) is a popular choice of companion animal, with an estimated population of 0.9 (PFMA, 2018) to 1 million (PDSA, 2018), making it the third most commonly kept mammalian species in the United Kingdom (PFMA, 2018). However, in their annual audit of welfare of companion animals in the UK in 2011, the People's Dispensary for Sick Animals (PDSA) identified the rabbit as one of the most neglected species (PDSA, 2011). Subsequent studies have confirmed a range of potential welfare issues (e.g. Rooney et al., 2013a) and a recent large-scale survey commissioned by the Royal Society for the Prevention of Cruelty to Animals (RSPCA) quantified (Rooney et al., 2014) and prioritised (Rooney et al., 2013a,) these, identifying fear of handling to be amongst the most important.

A study by Mullan and Main (2007) found that 20% of owners were not confident handling their rabbit(s). A larger survey found even higher levels of owner uncertainty (Rooney et al., 2014), with

27% of owners unwilling to classify themselves as “very confident”. This lack of confidence is likely to lead to poor, possibly unsafe handling practices that can exacerbate the rabbit’s fear response when handled. In fact, when observed in their own home, 61% of 221 rabbits were seen to show signs of fear when handled by their owners (Rooney et al., 2014). Common fear behaviors include crouching or hunching, freezing with the ears held back, running away when approached and aggressive behavior towards handlers (Magnus, 2005). Thumping of the hind limbs and growling or grunting are also signs that a rabbit is fearful (McBride, 2014). However, owners’ apparently often lack the ability to recognise that these behaviors can be symptomatic of fear; when asked in a questionnaire, 45.7% of owners reported their rabbit to be calm when handled, however an objective observer reported that only 33.8% were calm when handled by their owner (Rooney et al., 2014).

Despite the difficulty many owners have, handling is often necessary to facilitate veterinary visits, to perform routine health checks and to move rabbits between enclosures. Overall, 85.6% of rabbits are handled at least weekly and only a small number (2.2%) are never handled (Rooney et al 2014). In addition, traditionally, and even today, rabbits are often purchased as children’s pets, with surveys reporting 25% (PDSA, 2016), 39% (Edgar and Mullan, 2011), and even 45% (Mullan and Main, 2006) of rabbits being bought for a child. Young children are likely to handle rabbits frequently and sometimes inappropriately, hence it is unsurprising that 49% of rabbits are unable to be handled easily by children (Mullan and Main, 2007).

Fear is an aversive emotion, and when prolonged, repeated or intense, it can also have several long-term and potentially fatal consequences for rabbits. Cardiomyopathy, lymphopenia and reduced renal blood flow can all result, as well as gut stasis, a common condition for which stress is a known

cause (Varga, 2014). Hence fear of handling is a key area of welfare concern for which research is required to find methods of management and amelioration.

## **Use of Products to Reduce Fear in Companion Animals**

Non-pharmacological products that are marketed for reducing fear and anxiety are becoming increasingly popular in cats and dogs, when used alongside improved handling and behavioral training (Cracknell and Mills, 2008). Synthetic pheromone treatments and herbal remedies are commonly used ‘alternatives’ to pharmacological products. Efficacy data are usually lacking, so it is essential that the efficacy of such alternative treatments is tested. If they are ineffective they are at best a waste of client money and trust, but could also exacerbate problems and delay the use of more effective treatments.

Nutraceuticals (products derived from food sources and believed to have health or medical benefits) are growing in popularity and trials suggest that some may have potential value when used on ponies (McDonnell et al., 2013), horses (McDonnell, 2014), dogs (Beata et al., 2007a) and cats (Beata et al., 2007b). Interest has also grown in the use of natural herbs for reducing stress and anxiety in companion animals, including rabbits.

Valerian is an example of an herb which has the potential to improve animal welfare. It is derived from the root of the *Valeriana officinalis* plant (Hatteshol et al., 2008) and it has been shown to have anxiolytic effects on rodents and humans (Murphy et al., 2010, Becker et al., 2010). The suggested mode of action involves valerenic acid, the predominant anxiolytic root extract, interacting with gamma amino butyric acid (GABA) receptors (specifically GABA<sub>A</sub> receptors) in the central nervous

system (Murphy et al., 2010). This interaction causes neural inhibition by increasing the uptake of chloride ions, thus magnifying the hyperpolarisation effect of GABA on the neurons. Further neural inhibition could also be a result of other valerian root extracts inhibiting GABA transaminase activity, preventing the breakdown of GABA (Murphy et al., 2010).

Valerian has been shown to have relaxant properties for humans and is sometimes used in treatment of insomnia (Donath et al., 2000). Apparent reduced behavioral responsiveness could potentially be due to either an anxiolytic or a sedative effect. The mode of action is critical as the use of agents which induce sedation, with no true anxiolytic properties, is contra-indicated in animals showing signs of fear or anxiety. Other than preventing the animal from injuring itself in cases of extreme behavioral response, sedation alone is of limited benefit and may even make the problem worse; if it renders the animal unable to escape or utilise its natural coping mechanisms, thus potentially increasing the risk of sensitisation to the stressor. The animal will likely then be more (not less) fearful of the stimuli when it is next encountered. It is therefore vital to determine whether a product calms and hence facilitates reduced fear to stimuli or simply sedates and thus reduces avoidance responses.

Studies examining the effects of Valerian on vigilance in healthy human volunteers, suggest it does not result in sedation (Kuhlmann et al., 1999; Hallam et al., 2003; Gutierrez et al., 2004;). Studies using animal models similarly found Valerian extracts induced pronounced anxiolytic effects in rats with no associated reduction in locomotor activity, nor prolongation of ether-induced anaesthesia (Hattesoehl et al., 2008). Valerenic acid has also produced anxiolytic effects on mice with no effects on motor behavior or coordination (Murphy et al., 2010) and pigs exposed to a Valerian-containing herbal product during transportation had lower heart rates compared to controls (Peeters et al., 2004). There are currently several valerian-containing products, marketed as stress reduction aids,

139 primarily for cats and dogs, and available in varying forms including sprays, dermal spot-ons and  
140 products for oral administration.

141  
142 Pet Remedy® is a Valerian-based herbal product developed in the United Kingdom and  
143 manufactured and distributed through Unex Designs Ltd.. It is marketed as a safe, natural remedy  
144 for the treatment of stress and anxiety in all companion animals, including small mammals. The  
145 product range comprises a plug-in diffuser, battery-operated atomiser and a calming spray (Pet  
146 Remedy, 2017a). The spray is likely the most practical for rabbits, as many are kept outdoors. It is a  
147 pH-neutral, water-based formulation containing a blend of essential oils. Valerian oil is the principle  
148 component with smaller quantities of vetiver, basil and sage. Further constituents include  
149 polysorbate 20, a surfactant to facilitate mixing of the essential oils and the water base, and sodium  
150 benzoate and potassium sorbate as preservatives. Manufacturer guidelines suggest optimal results  
151 when applied to animal bedding, handler clothing or directly on to the animal's coat (Pet Remedy,  
152 2017a). Positive anecdotal reports abound, but to date, there have been a limited number of mainly  
153 unpublished trials into the efficacy of Pet Remedy for cats and dogs (Pet Remedy, 2017b; Barrington,  
154 2014; Taylor and Madden, 2016).

155  
156 These preliminary studies have produced contradictory and to our knowledge no such studies  
157 currently exist for rabbits. Fear of handling, as well as of novel environments and open spaces have  
158 been identified as key areas of welfare concern for this species (Rooney et al., 2014). Consequently,  
159 should Pet Remedy be effective at reducing fear in rabbits, it could have potential to significantly  
160 improve welfare. Therefore, this study seeks to test the efficacy of Pet Remedy on rabbits, whilst  
161 overcoming the limitations of previous studies on other species. Small group sizes (Barrington, 2014)  
162 and lack of observer blinding are two potential issues as, even in trials with placebo treatments, it is  
163 likely that experimenters could identify the active substance in Pet Remedy from its strong odour  
164 (Barrington, 2014; Taylor and Madden 2016). In the current study, we used a rehoming population



of rabbits to provide large group sizes, and a scented mask and nose clip were worn by the experimenter to mask the smell of the Pet Remedy and hence improve observer blinding. Since the subjects were of diverse and unknown histories, we expected wide ranging initial responses. The experimental design is within-subjects, with all rabbits undergoing both treatments (Pet Remedy and placebo) on successive days and responses compared to their own baseline levels.

The study tests the response of fifty rabbits when encountering an experimenter and a novel environment scented with Pet Remedy or a placebo substance of distilled water. The rabbits' responses towards a novel handler during an initial approach, when being picked up, handled and when placed in a novel arena were recorded. We measure behavioral responses but also take physiological measurements, i.e., heart and respiratory rate during handling. The test procedure utilised was adapted from that piloted by Rooney et al. (2014). We use a novel arena (Prut and Blezung, 2003) to allow measurement of its general confidence and its response to standardised behavior from a person to be measured.

We predict that if Pet Remedy has the effects claimed, individual rabbits will show calmer responses to handling, more positive, relaxed behaviors and fewer negative (fear and anxiety) behaviors after Pet Remedy as compared to placebo treatments. We hypothesise that significant decreases in heart and respiratory rates will be seen following Pet Remedy administration when compared to baseline and placebo levels. We also test whether Pet Remedy exerts a sedative effect on rabbits as locomotory behavior would be expected to decline if the action is sedative.

## **Methods and Materials**

188 *Ethical Approval*

189 This study was approved by the Animal Welfare and Ethical Review Board (AWERB) of the University  
190 of Bristol; Veterinary Investigation Number: VIN/16/001.

191 *Subjects*

192 Fifty rabbits were sourced from two locations; 14 rabbits from a privately-owned rehomed group in  
193 Taunton, and 36 from a rehoming centre in Gloucestershire. All testing was carried out in January  
194 and February 2016. Rabbits were minimally handled outside of the test procedures and so sex was  
195 recorded as reported by previous owners; 32 female and 18 male. All except three were neutered.  
196 The rabbits were of varying breed and were classified according to what they most closely  
197 resembled. The most common were Lionhead and their Crosses (n=9), Mini/Dwarf Lops and their  
198 crosses (n=8) and Cross Breeds (n=8). Rabbits were housed in various hutch and enclosure types; 18  
199 were kept in wooden hutches with no attached runs and 32 were housed in either large wooden  
200 sheds with access to an outdoor enclosure or indoor runs. Most rabbits were housed individually  
201 (n=22) or in pairs (n=23), three were housed in a group and the remaining two housed within a  
202 group of four. All rabbits remained in the care of their owner or rehoming centre after completion of  
203 the study.

204

205 Any rabbits with a pre-existing illness and those that were known to be aggressive towards handlers  
206 were excluded. All subjects had been at the location and in the same grouping for a minimum of one  
207 week. Where rabbits were kept in pairs, only one was used per cohort group. The second was tested  
208 at least a week later to avoid carry-over effects from approaching and picking up first rabbit. Where  
209 rabbits were housed in groups of three or more, two were used per cohort but both were subjected  
210 to the same treatment on the same day and there was a gap of at least one hour between rabbits.

To avoid unconscious bias (e.g., where calmer animals were selected first), the rabbit from each pair or group with the darkest fur or markings was used first.

### *Experimental design*

The design of this study was within- subjects so each rabbit was compared to itself because the sample was a mix of sexes and breeds. Each subject rabbit was tested three times by the same experimenter (SU) over three consecutive days. Baseline measures were taken on day 1 with no product being applied and on day 2 half the rabbits were randomly assigned to be exposed to Pet Remedy and half to placebo, by the experimenter who was blinded to the treatment. On day 3, the other substance was applied. The total population (n=50) was divided into six cohorts, each containing between six and ten rabbits which were tested over the same three-day period. For each cohort, one Pet Remedy (Unex Designs Ltd Manufactured 12/12/2015) and one placebo bottle were used. Twelve spray bottles were labelled one to twelve, (six Pet Remedy and six placebo containing distilled water) by an assistant, and randomly allocated as the first or second treatments to ensure the experimenter remained blind.

Each rabbit cohort was divided into two groups. Half (Group A) were tested in the morning and half (Group B) in the afternoon (Table 1). To avoid cross-contamination of the arena with the product, all rabbits tested in the morning received the same treatment, whilst all rabbits tested in the afternoon received the other. When the morning group had been completed, the arena was cleaned (using Beaphar® Deep Clean) and testing was stopped for 45 minutes to allow the arena to air out and eliminate residual smell for the later tests. At this point, the experimenter changed laboratory coat and the towel used to cover the carrier in which rabbits were moved.

*Table 1: Order of treatments received after Day 1 baseline by each cohort of rabbits*

Cohort	Group	Day Two Product	Day Three Product
<b>1</b>	a) n=4	Placebo	Pet Remedy
	b) n=4	Pet Remedy	Placebo
	Total n=8		
<b>2</b>	a) n=4	Placebo	Pet Remedy
	b) n=2	Pet Remedy	Placebo
	Total n=6		
<b>3</b>	a) n=4	Pet Remedy	Placebo
	b) n=4	Placebo	Pet Remedy
	Total n=8		
<b>4</b>	a) n=5	Placebo	Pet Remedy
	b) n=5	Pet Remedy	Placebo
	Total n=10		
<b>5</b>	a) n=5	Pet Remedy	Placebo
	b) n=4	Placebo	Pet Remedy
	Total n=9		
<b>6</b>	a) n=5	Pet Remedy	Placebo
	b) n=4	Placebo	Pet Remedy
	Total n=9		

#### *Experimental procedure*

In each rehoming centre the testing arena was assembled in a quiet, contained area out of olfactory contact with any of the subject animals. The wooden, collapsible arena, measuring 2m x 2m with 1m high walls was erected. Its floor was divided into nine equal squares for measurement of locomotory activity. A video camera was set up on a tripod on one side with a full view of the floor

244 and all tests filmed (Sony® Handycam DCR-SR58) in case behaviors were missed and for inter-  
245 observer testing.

246

247 The experimenter obtained basic demographics from the center owner, but did not make contact  
248 with any animals prior to the start of testing. For testing the experimenter wore a white long-  
249 sleeved, knee-length laboratory coat, white face mask scented with lemon and a nose clip to prevent  
250 distinction of the scent of Pet Remedy.

251

252 Prior to the start of testing on days 1 and 2, the appropriate treatment spray was applied to the  
253 experimenter's coat and a towel which was initially placed in a wire carrier used to transport the  
254 rabbit and then placed into the novel arena with the rabbit. One spray (approximately 0.2 ml) was  
255 applied to each of the experimenter's cuffs, one to the body of the coat and three sprays to the  
256 towel (1.2ml in total).

257

## 258 ***Testing Protocol***

259 Rabbits were tested in the same order and at approximately the same time of day on all three days.  
260 If any rabbit showed extreme negative behaviors at any point during a test (e.g., open-mouth  
261 breathing), the test would be stopped immediately, and the rabbit returned to its home enclosure.  
262 This never occurred.

263 The testing protocol had seven sub elements:

264 ***1. Hutch approach:*** The experimenter placed a small animal wire carrier close to the hutch or  
265 enclosure but out of sight of the rabbit. She approached and placed her hand against the hutch bars  
266 or opened the enclosure door (if there were no bars/wire) and held her hand out, at least one body

length from the rabbit. She remained for thirty seconds and the rabbit's initial and maximum (most positive) responses were recorded as well as their position for the majority of the time period (Table 2).

**2. Pick up:** The experimenter opened the hutch door or entered the enclosure and started the stopwatch. She allowed the rabbit to approach before moving to capture it calmly. The rabbit was carried close to the ground and placed in the wire carrier, lined with the towel. The time taken to pick up and the rabbit's behavioral responses (positive and negative) were recorded.

Once the rabbit was placed in the wire carrier, a large towel was placed over it and it was moved to the novel arena. To standardise exposure to the product, all rabbits remained in the carrier between one and two minutes.

**3. Novel Arena** The video camera was activated; the towel was removed from the wire carrier and the carrier lifted into the arena. The rabbit was lifted out of the carrier and placed in the central square of the arena, the carrier removed and the towel placed close to the centre of the arena. Latency to move, number of squares passed into and interactions with the towel, rearing frequency and the number and types of positive (exploring, grooming, rearing, lying stretching out, sniffing) and negative behaviors exhibited (freeze, scratch corners, escape attempts) were then recorded over a two minute period.

**4. Experimenter hand and carrot:** The experimenter placed her hand, holding a piece of carrot in the arena corner, at least one square away from the rabbit. The closest the rabbit moved towards the carrot within a 30 second period was recorded.

**5. Experimenter in novel arena:** The experimenter quietly entered the arena and sat for 30 seconds, cross-legged. The rabbit's response, the frequency of positive and negative behaviors was recorded.

**6. Experimenter handling:** The experimenter placed the towel on her crossed legs before moving to picked up the rabbit and placed it on the towel. She then spent one minute gently stroking the rabbit

from head to tail and assessing Body Condition Score (BCS; Mullan and Main, 2006). During the second minute, respiratory rate and heart rate were measured, subjective ratings were recorded for overall level of reactivity during handling and level of restraint required to maintain hold of the rabbit. Number of escape attempts were also recorded.

**7. Physiological measures:** The experimenter observed the rise and fall of the chest to count respiratory rate, and placed a stethoscope to determine heart rate, counted over 15 seconds and converted to breaths/beats per minute.

After completion of the test, recording was stopped, and the rabbit returned to the wire carrier and moved back to their home enclosure. The walls and floor of the novel arena and the wire carrier were sprayed with low-odour disinfectant (Beaphar® Deep Clean) and wiped down. The experimenter disinfected her hands with hand gel of minimal scent (Cuticura® Original) before testing the next rabbit.

### *Statistical Analysis*

Statistical analysis was performed using IBM® SPSS® Statistics 23 for Windows. Initially 113 variables were recorded during testing and descriptive analysis and histograms determined the spread for each variable. Variables describing very rare behaviors were eliminated, as were those showing very little variation. Where appropriate, frequencies were scaled to improve spread, and binary responses for specific subtests were grouped over the entire test, thereby producing 20 meaningful variables for subsequent analysis. Within each subtest, we compared the variables using a Spearman Rank Correlation test. For any correlations above 0.7 (i.e., over 50% of variation being explained), the more subjective of the measures was eliminated. Thus, from the Experimenter Handling sub-test

314 we eliminated 'Restraint required during handling' as this was highly correlated to 'Number of  
315 escape attempts' (0.773) leaving 19 variables (Table 2).

316

317 A second trained observer blindly analysed 18 of the testing sessions. Her measures for seven  
318 variables were chosen including at least one from each subtest, the variable showing the most  
319 variability between rabbits. Observers were compared for inter-observer reliability using Kendall's  
320 Index of Concordance and Chi squared tests for continuous and binary variables respectively.

321



323 *Table 2: Description of the 20 variables derived from data recorded during seven test components*

Test component	Variable	Description	Scale
1. Hutch Approach	Positivity of Response (At 0 secs)	Rabbit's initial response to experimenter's hand at front of hutch or in doorway of enclosure	<b>0</b> - Freeze/Out of sight <b>1</b> - No Response, Withdraw, Out of sight <b>2</b> - Turn head, no other movement, Come out of hiding <b>3</b> – Approach, Sniff hand <b>4</b> - Contact hand
	Maximum Positivity of Response	Rabbit's maximum positive response over 30 seconds to experimenter's hand at front of hutch or in doorway of enclosure	<b>0</b> – Freeze, Out of sight <b>1</b> - No Response, Withdraw, Out of sight <b>2</b> - Turn head, no other movement, Come out of hiding <b>3</b> – Approach, Sniff hand <b>4</b> - Contact hand
2. Pick up	Latency to Pick Up	Time taken (seconds) to capture rabbit in hutch or enclosure	<i>Grouped for analysis:</i> <b>1</b> - <10 secs <b>2</b> - <30 secs <b>3</b> - <60 secs <b>4</b> - <120 secs <b>5</b> - 120+ secs

	Negative Responses to Being Picked up	Number of negative responses observed during capture of rabbit	<b><i>Negative responses:</i></b> Run Away Thump Vocalise Freeze Struggle Slightly Struggle Intensely Bite/Scratch/Kick
3. Novel Arena	Latency to Move	Time taken (seconds) to move after being placed into novel arena	<b><i>Grouped for analysis:</i></b> <b>1</b> - 0 secs <b>2</b> - 1-5 secs <b>3</b> - 6-10 secs <b>4</b> - 11+ secs
	Number of Squares Entered	Number of squares moved into with all feet during two-minute test period	
	Number of Interactions with Towel	Number of interactions with the towel during two-minute test period	
	Number of Rears	Number of rears observed during two-minute test period	<b><i>Grouped for analysis:</i></b> <b>1</b> - 0 rears <b>2</b> - 1-5 rears <b>3</b> - 6-10 rears <b>4</b> - 11+ rears
	Number of Different Negative Behaviors	Number of types of negative behaviors observed during two-minute novel arena test	<b><i>Negative behaviors:</i></b> Freeze Scratch Corners Frantic to Get Out

		Number of Different Positive Behaviors	Number of types of positive behaviors observed during two-minute novel arena test	<b>Positive behaviors:</b> Explore Groom Rear Up Lie Stretched Sit and Sniff
		Positivity of Response	Most common behavior observed during the two-minute novel arena test	<b>0</b> - Frantic to get out <b>1</b> - Freeze/Scratch Corners <b>2</b> - Sit and Sniff <b>3</b> - Explore/Rear Up <b>4</b> - Groom <b>5</b> - Lie Stretched
4.	Experimenter hand and carrot	Closeness to Experimenter Hand	A 3-point scale of minimum distance between experimenter's hand and the rabbit over 30 seconds	<b>1</b> - Furthest corner away, Two squares away <b>2</b> - One square away, Same square <b>3</b> - Contact, Eats Carrot
5.	Experimenter in novel arena	Number of Different Negative Behaviors Observed	Number of different negative responses observed after experimenter sat in novel arena with rabbit. Each behavior was counted only once	<b>Negative Behaviors:</b> Freeze Attempt Escape Thump Vocalise
		Number of Different Positive Behaviors Observed	Number of positive behaviors observed during 30 second period in which	<b>Positive Behaviors:</b> Approach Sniff or Contact

		experimenter sat in arena with rabbit. Each behavior was counted only once	Sit but alert Carry on as before Climb on lap
6. Experimenter handling	Positivity of Response	Scale of positivity of behavioral responses to experimenter handling of rabbit.	<b>0</b> - Can't be handled <b>1</b> - Bites/Scratches/Kicks, Struggles Intensely <b>2</b> - Freezes, Struggles Slightly <b>3</b> - Tense, no struggling <b>4</b> - Calm
	Number of Escape Attempts	Number of attempts to escape during two-minute experimenter handling	
	Reactivity during Handling	Subjective scale of how reactive experimenter perceived rabbit to be during two-minute handling period	<i>4-point scale where:</i> <b>1</b> - Very calm <b>4</b> - Not calm at all
	Restraint Required During Handling	Scale of the level of restraint required by experimenter to maintain hold on rabbit during two-minute handling test	<b>1</b> - Sits unheld <b>2</b> - Held loosely <b>3</b> - Held firmly <b>4</b> - Held tightly
7. Physiological Measures	Heart Rate	Heart rate of rabbit at end of all tests	Measured over 15 second and calculated per minute 60 seconds

Respiratory Rate	Respiratory rate of rabbit	Measured over 15
	at end of all tests	second and calculated
		per minute 60 seconds

324

325 Data Reduction using Principal Components Analysis was trialled on the data but failed to reduce the  
 326 variables to a smaller number of meaningful factors. Therefore, the raw variables were used in  
 327 subsequent analysis and results interpreted with caution due to the dangers of multiple testing.

328

329 We examined the effect of treatment (Pet Remedy or Placebo), the order in which treatments were  
 330 administered and the individual rabbit's baseline response (included as a covariate), and the  
 331 interaction between order and treatment on each variable (Table 3).

## 332 **Results**

333 Of the seven variables tested for inter-observer reliability, all showed significant agreement between  
 334 the two observers ( $p < 0.005$ ).

335

336

337 *Table 3: Results of Repeated Measures General Linear Model analysis of all 19 variables, including F-*  
 338 *and P- values for each. P<0.05 (\*) was accepted as a significant result.*

Variable		ANOVA Test Results							
	Name	Treatment		Baseline		Study Day (Order)		Treatment x Order Interaction	
		F	P	F	P	F	P	F	P
1. Hutch	Positivity of	0.98	0.327	2.85	0.007*	1.05	0.311	0.00	0.984
Approach	Response (at 0 seconds)								
	Maximum	0.38	0.539	2.33	0.024*	1.00	0.324	0.14	0.706
	Positivity of Response								
2. Pick up	Latency to Pick	4.08	0.049	4.43	<0.001*	0.01	0.906	0.69	0.410
	up		*						
	Negative	0.14	0.706	4.38	<0.001*	0.67	0.417	0.38	0.539
	Responses to being picked up								
3. Novel	Latency to	0.08	0.782	2.88	0.006*	0.00	0.961	2.39	0.129
Arena	Move								
	Number of	0.02	0.902	5.54	<0.001*	0.34	0.560	3.63	0.063
	Squares Entered								
	Number of	0.10	0.748	3.40	0.001*	0.03	0.859	2.60	0.114
	Interactions with Towel								

	Number of Rears	2.67	0.109	2.68	0.001*	6.91	0.012	0.30	0.589
							*		
	Number of Different Negative Behaviors	0.00	0.983	2.59	<0.013*	1.50	0.226	1.12	0.295
	Number of Different Positive Behaviors	4.52	0.039	2.63	0.012*	1.42	0.240	0.79	0.380
			*						
	Positivity of Response	0.12	0.729	4.49	<0.001*	0.01	0.938	2.92	0.094
4.	Closeness to Experimenter Hand with Carrot	0.44	0.511	3.60	0.001*	0.01	0.938	1.02	0.318
5.	Number of Negative Behaviors Observed	2.97	0.092	3.67	<0.001*	2.71	0.107	0.00	1.000
	Number of Positive Behaviors Observed	1.11	0.296	2.69	0.010*	1.68	0.201	1.95	0.169
6.	Positivity of Response	2.09	0.155	2.44	0.019*	0.09	0.767	0.11	0.737

Handling	Number of	1.72	0.196	2.73	0.009*	2.57	0.116	8.04	0.007*
	escape								
	attempts								
	Reactivity	0.05	0.825	2.44	0.019*	0.52	0.473	7.93	0.007*
	during								
	handling								
7.	Heart Rate	4.41	0.04*	3.37	0.002*	0.14	0.710	0.20	0.657
Physiological	Respiratory	0.81	0.373	6.87	<0.001*	0.97	0.329	0.02	0.900
Measures	Rate								

339

340 Three variables were significantly affected by treatment (Table 3); Latency to Pick up), Number of  
341 Different Positive Behaviors in the Novel Arena and Heart Rate Latency to Pick up decreased with  
342 both Placebo and Pet Remedy treatments, but the greatest reduction relative to baseline  
343 (M=3.80±0.95) was seen with Placebo (M=3.46±0.91) as compared to Pet Remedy Treatment  
344 (M=3.78±0.97; Figure 1A; Table 4).

345

346 The Number of Different Positive Behaviors observed in the Novel Arena was approximately the  
347 same with Placebo (M=1.98±0.92) as at Baseline (M=1.98±0.77) but significantly higher following  
348 treatment with Pet Remedy (2.388±1.07; Figure 1B).

349

350 Heart Rate was marginally increased during Placebo trials (M=265.23±29.56) as compared to  
351 baseline (M=263.67±27.77) whilst Treatment with Pet Remedy was accompanied by a decrease in  
352 heart rate (M=253.23±33.70; Figure 1C). *Table 4: Mean and Standard Deviation for all variables*  
353 *shaded are the variables significantly affected by treated*

<u>Variable Name</u>	<u>Condition</u>
----------------------	------------------



		<u>Baseline</u>		<u>Placebo</u>		<u>Pet Remedy</u>	
		Mean	Sd	Mean	Sd	Mean	Sd
1.	Positivity of	1.80	0.904	1.68	0.819	1.55	0.792
Hutch	Response (at 0						
Approach	seconds)						
	Maximum	2.22	0.932	2.04	1.009	1.94	0.944
	Positivity of						
	Response						
2.	Latency to Pick up	3.80	0.948	3.46	0.908	3.78	0.872
Pick up	Negative	1.90	1.129	1.98	1.000	1.94	1.162
	Responses to						
	being picked up						
3.	Latency to Move	1.92	0.853	1.58	0.810	1.61	0.812
Novel Arena	Number of	16.80	10.882	11.12	9.506	11.04	7.681
	Squares Entered						
	Number of	3.38	1.999	2.41	1.645	2.49	1.781
	Interactions with						
	Towel						
	Number of Rears	1.90	0.707	2.08	1.017	2.27	1.036
	Number of	1.34	0.982	1.20	1.06	1.55	1.308
	Different						
	Negative						
	Behaviors						
	Number of	1.98	0.769	1.98	0.92	2.39	1.077
	Different Positive						
	Responses						
	Positivity of	2.39	0.45	2.31	0.56	2.30	0.506
	Response						

4.	Closeness to	2.04	0.781	1.62	0.81	1.71	0.842
Experimenter	experimenter						
Hand and	hand						
Carrot							
5.	Number of	0.84	1.595	1.02	1.65	0.59	1.606
Experimenter	Different						
in Novel Arena	Negative						
	Behaviors						
	Observed						
	Number of	1.38	1.123	1.56	1.18	1.265	1.186
	Different Positive						
	Behaviors						
	Observed						
6.	Positivity of	2.38	0.780	2.54	0.91	2.27	0.818
	Response						
Experimenter	Number of	1.12	1.35	0.92	0.99	1.16	1.179
Handling	Escape Attempts						
	Reactivity during	3.82	0.77	3.74	0.78	3.73	0.758
	handling						
7.	Heart Rate	263.67	27.77	265.23	29.56	253.23	33.70
Physiological	Respiratory Rate	144.76	32.63	144.20	36.74	149.06	39.84
Measures							

354

355 [ FIGURE 1 (A,B,C) ]

All variables analysed were significantly affected by an individual rabbit's baseline data, seven variables with P-values of less than 0.001 (Table 3), indicating an individual's response during the baseline test was highly predictive of how they would respond in subsequent tests.

Study Day (Day 2 or Day 3) had a significant effect on only one variable, the number of rears observed during the novel arena test ( $F_{(1,45)}=6.91, P=0.012$ ). There was a significant increase in the number of rears observed from Day 2 ( $M=2.12\pm1.04$ ) to Day 3 ( $M=2.30\pm1.02$ ) (Figure 2).

## **[FIGURE 2]**

The interaction between order and treatment was significant for two variables; Number of Escape Attempts during Experimenter Handling ( $F_{(1,47)}=8.04, P=0.007$ ) and Reactivity during Handling ( $F_{(1,47)}=7.93, P=0.007$ ; Table 3). In those rabbits treated with Placebo first, the Number of Escape Attempts decreased from Baseline ( $M=1.12\pm1.35$ ) to Day 2 ( $M=0.92\pm1.00$ ). When these rabbits were then exposed to Pet Remedy on Day 3, mean number of escapes decreased further ( $M=0.71\pm1.08$ ). In contrast in those that received Pet Remedy first, the Number of Escapes Attempts increased from Baseline to Day 2 ( $M=1.60\pm1.12$ ), but decreased greatly in relation to both baseline and Day 2 data when treated with placebo on Day 3 ( $M=0.92\pm1.00$ ; Figure 3A).

Both groups of rabbits were deemed more Reactive on Day 2 than Day 1, irrespective of product used. Those that received Pet Remedy on Day 2 were more reactive during both treatment tests than those who received Placebo first. A decrease in Reactivity from Day 2 to 3 was demonstrated by both groups, both groups falling below baseline data on this day (Figure 3B).

[FIGURE 3 A,B]

## Discussion

This study, overall, saw few differences in behavior between trials in which placebo and Pet Remedy were administered. Of the 19 variables analysed, only three were significantly different according to treatment. Treatment with Pet Remedy was associated with a significant decrease in Heart Rate during handling and a significant increase in the Number of Different Positive Behaviors, observed in a novel environment compared to placebo. Although the Latency to Pick up the rabbit was lower in trials with placebo, since this was the first encounter with the experimenter during the test and often followed a Pet Remedy trial, this too may support a calming effect. In contrast, other variables such as Respiratory Rate, Reactivity to handling and Number of Different Negative Behaviors Observed in a Novel Arena or during Handling showed no significant differences with treatment. Two variables, Number of Escape Attempts and the subjective rating for Reactivity During Handling showed complex treatment/order interactions which require further investigation.

A previous study of oral administration of valerian, the main active component of Pet Remedy, produced a decrease in heart rate in pigs during simulated transport (Peeters et al., 2004). Despite the difference in method of administration in the current trial (olfactory rather than oral), Pet Remedy was again associated with a significant decrease in heart rate when compared with baseline and placebo in rabbits. However, heart rates were relatively high throughout this study averaging 260 compared to normal resting levels, which are 154-300 (Varga, 2014). These high heart rates may partly be explained by the exercise during capture, which occurred several minutes before testing,

but we suggest it predominantly shows that handling, and in particular handling by a novel person, is particularly stressful to rabbits. Heart rate is reduced by Pet Remedy administration and it is possible that this decrease was associated with a concurrent reduction in stress, (the concurrent increase in positive behaviors suggests that this was not a reduction in positive arousal), but the Pet Remedy treatment alone is insufficient to reduce heart rate to the low end of the normal range. It is interesting to note that no concurrent significant change in respiratory rate was seen. As respiratory and heart rates are usually closely linked, a change would have been expected. However, respiratory rates were very high, averaging 146, compared to normal 30-60 breaths per minute (Varga, 2014), suggesting an extreme physiological response that may have been insensitive to relatively minor treatment differences.

Treatment with Pet Remedy was also associated with a significant increase in the Number of Different Positive Behaviors observed in the Novel Arena test. Positive behaviors such as exploring, grooming, rearing up, lying down, sitting and sniffing the environment are unlikely to be seen when rabbits are distressed and their increase after Pet Remedy administration supports the hypothesis that Pet Remedy has a calming effect. Other behaviors measured during handling and hutch approaches which may also be indicators of relaxation or distress were not significantly affected. However, the increase in the variety of positive behaviors, combined with no significant difference in the Number of Squares Entered, suggests that Pet Remedy did not cause sedation.

The time taken to pick up the rabbits from their home enclosure decreased from baseline with both treatments, but a greater effect was seen with the placebo than with Pet Remedy, although this is only marginally significant ( $P=0.049$ ). This may seem unexpected if the Pet Remedy is calming, one may expect the rabbit to be easier to pick up. However, one has to consider the order of the test components. Capture occurred immediately after the initial hutch approach when the rabbits had

had very little exposure to the odour. Hence, they were likely more affected by their previous interaction with the experimenter than the current trial. The rabbits when treated with Pet Remedy had experienced either a baseline or a placebo treatment on the previous day, (whilst placebo followed baseline or Pet Remedy) , so differences in capture time may have a carry-over from the last handling session. This points to the need to investigate the effect of Pet Remedy administered over successive handling and testing sessions to test whether it aids habituation or systematic desensitization and avoid sensitization in the animals.

The Number of Escape Attempts and the subjective rating of Reactivity during Handling were both significantly affected by the order in which the two treatments were given over the three days. Rabbits that were treated with Pet Remedy on Day 2 were rated as more Reactive to the experimenter handling, than those who received placebo on that day. This same group were again later classified as more Reactive on Day 3 when they were treated with the placebo spray. Escape attempts increased on Day 2 in rabbits given Pet Remedy first and subsequently decreased while those getting placebo first experienced a gradual increase in escape attempts. It is possible that escape attempts only happen when the animal is not freezing with fear and hence sufficiently relaxed to explore its environment, or they may be indicative of fear of the arena. The exact cause of this order dependent, behavioral change is unclear, but it does support the conclusion that rabbits were not sedated when exposed to Pet Remedy.

Although heart rate was reduced none of the behavioral results support the possibility that rabbits exposed to Pet Remedy were sedated. There was no significant difference in locomotory behavior recorded between Pet Remedy and placebo trials, nor was there any evidence that rabbits were more wary or fearful of the experimenter on the trials following Pet Remedy administration. In fact, the number of rears observed in the novel arena increased significantly from Day 1 through to Day 3,

in Pet Remedy as well as placebo conditions. This is likely due to habituation to the arena; with repeated exposure, the rabbits became more familiar and reared up to survey their surroundings. This would not be expected if the rabbits were sedated. This study however was only conducted over three days and we suggest that longitudinal studies involving repetitive handling of rabbits over longer periods of time are required with an emphasis on monitoring the animals for any signs of sedation as this could result in an inability to physically retreat from aversive and threatening stimuli.

The current study had several limitations. Although, we aimed to mask the smell of the Pet Remedy spray using a nose clip and lemon oil, and inter-observer reliability checks confirmed consistent, unbiased recording, there remains the possibility that the measures were insufficient to fully blind the tester or that the rabbits responded to the smell. It would therefore be valuable in future trials to be able to produce a placebo which smells similar but lacks the active ingredients of Pet Remedy.

Initial data reduction efforts failed to reduce the variables to meaningful underlying factors, therefore analysis was done on 19 raw variables. This raises the issue of repeat testing and plausible Type II errors and so individual results need to be interpreted with caution. The number of significant results is few, given the number of variables tested, although more than would be expected by chance alone. The results suggest that Pet Remedy may have some positive effect at reducing acute stress in rabbits and shows no evidence of sedating the rabbits, suggesting it may be of use in veterinary practice where visits are often short but can be particularly stressful, although application in the environment prior to handling is likely required. As this trial was only carried out over three days, the effectiveness of long-term usage remains to be tested.

476

477 For all the variables tested, an individual's response in the preliminary baseline test had a very  
478 significant effect on how they reacted in later tests. Variation in individual responses to stress within  
479 a population is widely acknowledged (Koolhaas et al., 1999) and would be expected to be a  
480 significant factor especially in a rehoming population where most subjects' histories are unknown  
481 and likely vary greatly. The within-subjects design and the use of baseline data as a covariate in the  
482 analysis meant that such inter-rabbit differences were taken into consideration. However, it is  
483 possible that rabbits of initially different reactivity, respond differently to Pet Remedy which may  
484 make significant effects more difficult to detect than in a more standardised population of animals.  
485 A rehoming population, however represents an important part of the pet population at which this  
486 product is aimed and hence this is a meaningful first study.

487

488

489 Apparent throughout this study, was the fact that within a population of fifty rabbits, the vast  
490 majority showed aversion to human approaches and handling. Very few voluntarily approached a  
491 person, and most struggled and attempted to escape from handling. Fear of handling is very  
492 common in the general population of pet rabbits (Mullan and Main, 2006). Mitigation via early  
493 socialisation, appropriate introduction to handling and optimal handling techniques are vital.

494

495 Early handling of rabbits is widely recognised as an effective method of preventing fear during  
496 handling later in life (Magnus, 2005; McBride, 2014). Research on laboratory populations shows  
497 handling kittens within the first week of life significantly affects their later behavioral responses  
498 (Bilko et al., 2000; Zucca et al., 2012) and handling around the time of nursing reduces fearfulness as  
499 an adult (Bilko et al., 2000; Pongrácz and Altbäcker, 2003). The technique used to handle rabbits is



also important. Wild rabbits are prey species, for whom being lifted off the ground signifies likely risk of death by predation. Since domestication has changed rabbits' natural behavior very little (Lehman, 1991). Lifting is also aversive to domestic rabbits. Handling them on the ground is generally acknowledged to be less aversive.

For those rabbits with established fears, behavior modification, incorporating techniques such as systematic desensitisation (DS), and counter-conditioning (CC) is recommended as an effective method by which to reduce fearful behaviors (Magnus, 2005). These techniques involve changing the animal's perception of the fear-eliciting stimulus, using a controlled gradual exposure (DS), whilst associating it with something positive, such as food (CC) and are well established in a number of species (e.g. Levine et al., 2007). Future validation and publication of optimal handling protocols (for adults and kittens), and behavior modification techniques have the potential to improve rabbit welfare further.

## Conclusions

We used a standardized test of rabbit-human behavior and analyzed rabbits' responses to a novel standardized handling test. This testing protocol produced a variety of measures which showed wide variation and high inter-observer reliability. Individual variation was great across the population, with baseline levels exerting a very pronounced effect on all aspects of behavior and physiology in later tests. This finding suggests the test was successful in measuring individual differences in behavioral responses.

The study also highlighted the high level of fear of handling within the rabbit population. Pet Remedy used on a handler's clothing and in a novel environment was associated with a significant

decrease in heart rate during handling and an increase in positive behaviors observed in a novel environment but produced no noticeable change in the rabbit's behavior towards the experimenter during a single administration. Use of Pet Remedy was also associated with a longer latency to pick up the rabbit, although this was possibly due to carryover effects from the previous testing session. Although there were no differences for most of the behavior measures, the reduction in Heart Rate and increase in positive behaviours suggest that Pet Remedy may have some potential value for rabbits during periods of acute stress and may thus be useful during veterinary visits and during initial handling and is worthy of further investigation.

### **Acknowledgements**

The authors wish to thank Martyn Barklett-Judge Managing Director of Unex Designs Ltd for funding student travel, providing products and funding production of this manuscript. We thank Alice Chamberlain at Windwhistle Warren Rabbit & Guinea Pig Rescue and Shirley Hughes for use of rabbits. Thank you also to Rae Todd of the Rabbit Welfare Association Fund for continued support, help with piloting and friendly advice, Richard Parker and Toby Knowles for statistical advice and Katherine McIndoe for invaluable help with inter-observer reliability and reference checking.

### **Competing interests:**

We declare that none of the authors have any competing interests

## References

- Barrington, E. A., 2014. study to assess the ability of 'Pet Remedy' to reduce stress in rescue cats (*Felis catus*). Unpublished BSc (Hons) Dissertation. University of Exeter.
- Beata, C., Beaumont-Graff, E., Diaz, C., Marion, M., Massal, N., Marlois, N., Muller, G., Lefranc, C., 2007. Effects of alpha-casozepine (Zylkene) versus selegiline hydrochloride (Selgian, Anipryl) on anxiety disorders in dogs. *J. Vet. Behav.: Clin. Appl. Res.* 2 (5), 175-183.
- Beata, C., Beaumont-Graff, E., Coll, V., Cordel, J., Marion, M., Massal, N., Marlois, N., Tauzin, J., 2007. Effect of alpha-casozepine (Zylkene) on anxiety in cats *J. Vet. Behav.: Clin. Appl. Res.* 2 (2), 40-46.
- Becker, A., Felgentreff, F., Schröder, H., Meier, B., Brattström, A., 2017. The anxiolytic effects of a Valerian extract is based on Valerenic acid. *BMC Complement Altern Med.* 14 (1), 267.
- Bilkó, Á., Altbäcker, V., 2000. Regular handling early in the nursing period eliminates fear responses toward human beings in wild and domestic rabbits. *Devel.Psychobiol.* 36, 78-87.
- Cracknell, N.R., Mills, D.S., 2008. A double-blind placebo-controlled study into the efficacy of a homeopathic remedy for fear of firework noises in the dog (*Canis familiaris*). *Vet. J.* 177, 80-88.
- Donath, F., Quispe, S., Diefenbach, K., Maurer, A., Fietze, I., Roots, I., 2000. Critical Evaluation of the Effect of Valerian Extract on Sleep Structure and Sleep Quality. *Pharmacopsychiatry.* 33 (2), 47-53.
- Edgar, J.L., Mullan, S.M., 2011. Knowledge and attitudes of 52 UK pet rabbit owners at the point of sale. *Vet. Rec.* 168, 353.
- Gutierrez, S., Ang-Lee, M., Walker, D., Zacny, J., 2004. Assessing subjective and psychomotor effects of the herbal medication valerian in healthy volunteers. *Pharmacol. Biochem. Behav.* 78 (1), 57-64.
- Hallam, K., Olver, J., McGrath, C., Norman, T., 2003. Comparative cognitive and psychomotor effects of single doses of *Valeriana officianalis* and triazolam in healthy volunteers. *Hum, Psychopharm. Clin.* 18 (8), 619-625.

569 Hatteshol, M., Feistel, B., Sievers, H., Lehnfeld, R., Hegger, M., Winterhoff, H., 2008. Extracts of  
 570 Valeriana officinalis L. s.l. show anxiolytic and antidepressant effects but neither sedative nor  
 571 myorelaxant properties. *Phytomedicine*. 15, 2-15.

572 Koolhaas, J.M., Korte, S.M., De Boer, S.F., Van Der Vegt, B.J., Van Reenen, C.G., Hopster, H., De Jong,  
 573 I.C, Ruis, M.A and Blockhuis, H.J., 1999. Coping styles in animals: current status in behavior and  
 574 stress-physiology. *Neurosci. Biobehav.* 23, 925-935.

575 Kuhlmann, J., Berger, W., Podzuweit, H., Schmidt, U., 1999. The Influence of Valerian Treatment on  
 576 'Reaction Time, Alertness and Concentration' in Volunteers. *Pharmacopsychiatry*. 32 (6), 235-241.

577 Lehmann, M., 1991. Social Behavior in young domestic rabbits under semi-natural conditions. *Appl.*  
 578 *Anim. Behav. Sci.* 32, 269-292.

579 Levine, E., Ramos, D., Mills, D.S., 2007. A prospective study of two self-help CD based desensitization  
 580 and counter-conditioning programmes with the use of Dog Appeasing Pheromone for the treatment  
 581 of firework fears in dogs (*Canis familiaris*). *Appl. Anim. Behav. Sci.* 105(4), 311-329.

582 McBride, E.A., 2017. Normal behavior and behavior problems. In: Meredith M, Lord B, editors.  
 583 BSAVA Manual of Rabbit Medicine. 1<sup>st</sup> ed. Gloucester: British Small Animal Veterinary Association; .  
 584 pp45-58.

585 McDonnell, S., Miller, J., Vaala, W., 2013. Calming Benefit of Short-Term Alpha-Casozepine  
 586 Supplementation During Acclimation to Domestic Environment and Basic Ground Training of Adult  
 587 Semi-Feral Ponies. *J. Equine Vet. Sci.* 33 (2), 101-106.

588 McDonnell, S., Miller, J., Vaala, W., 2014. Modestly Improved Compliance and Apparent Comfort of  
 589 Horses With Aversions to Mildly Aversive Routine Health Care Procedures Following Short-Term  
 590 Alpha-Casozepine Supplementation. *J. Equine Vet. Sci.* 34 (8), 1016-1020.

591 Magnus, E., 2005. Behavior of the pet rabbit: what is normal and why do problems develop? In  
 592 *Practice*. 27, 531-535.

593 Murphy, K., Kubin, Z.J., Shepherd, J.N., Ettinger, R.H., 2010. Valeriana officinalis root extracts have  
 594 potent anxiolytic effects in laboratory rats. Phytomedicine. 17, 674-678.

595 Mullan, S.M., Main, D.C. 2006. Survey of the husbandry, health and welfare of 102 pet rabbits. Vet.  
 596 Rec. 159, 103-109.

597 Mullan, S.M., Main, D.C., 2007. Behavior and personality of pet rabbits and their interactions with  
 598 their owners. Vet. Rec. 160, 516-520.

599 PDSA. (Peoples' Dispensary for Sick Animals) 2018. Peoples' Dispensary for Sick Animals Animal  
 600 Wellbeing Re.port [https://www.pdsa.org.uk/get-involved/our-campaigns/pdsa-animal-wellbeing-](https://www.pdsa.org.uk/get-involved/our-campaigns/pdsa-animal-wellbeing-report)  
 601 report 2018

602 Peeters, E., Driessen, R., Steegmans, R., Henot, D., Geers, R., 2004. Effect of supplemental  
 603 tryptophan, vitamin E, and a herbal product on responses by pigs to vibration. J. Anim. Sci. 82 (8),  
 604 2410-2420.

605 PFMA (Pet Food Manufacturers' Association) 2018. Pet population 2018  
 606 <https://www.pfma.org.uk/pet-population-2018> 6/7/2019

607 Pet Remedy How Pet Remedy Works. 2017a <http://petremedy.co.uk/how-to-use/> accessed  
 608 6/7/2019.

609 Pet Remedy. Clinical Trials. <http://www.petremedy.co.uk/clinical-trials> accessed 6/7/2019

610 Pongrácz, P., Altbäcker, V., 2003. Arousal, but not nursing, is necessary to elicit a decreased fear  
 611 reaction toward humans in rabbit (*Oryctolagus cuniculus*) pups. Dev. Psychobiol. 43, 192-199.

612 Prut, L., Blezung, C., 2003. The open field as a paradigm to measure the effects of drugs on anxiety-  
 613 like behaviors: a review. Eur. J. Pharmacol. 463, 3-33.

614 Rooney, N.J., Mullan, S., Blackwell, E.J., Saunders, R., Held, S. 2013a. Prioritising target issues in a  
 615 pet rabbit welfare campaign. Poster Presentation at Universities Federation for Animal Welfare

616 (UFAW), Barcelona. Available from:[http://www.ufaw.org.uk/downloads/events/ufaw-2013-](http://www.ufaw.org.uk/downloads/events/ufaw-2013-symposium-abstract-booklet-final-v3.pdf)  
617 [symposium-abstract-booklet-final-v3.pdf](http://www.ufaw.org.uk/downloads/events/ufaw-2013-symposium-abstract-booklet-final-v3.pdf)

618 Rooney, N.J., Held, S., Mullan, S., Blackwell, E.J., and Saunders, R., 2013b. The current state of  
619 welfare, housing and husbandry of the English pet rabbit population. Report to RSPCA.

620 Rooney, N.J., Blackwell, E.J., Mullan, S.M., Saunders, R., Baker, P.E., Hill, J.M., Sealey, C.E., Turner,  
621 M.J., Held, S.D.E., 2014. The current state of welfare, housing and husbandry of the English pet  
622 rabbit population. BMC Res Notes. 7, 942.

623 Taylor, S., Madden, J., 2016. The Effect of Pet Remedy on the Behavior of the Domestic Dog (*Canis*  
624 *familiaris*). Animals. 6 (11), 64.

625 Varga, M., 2014. The rabbit-friendly practice. In: Meredith M, Lord B, editors. BSAVA Manual of  
626 Rabbit Medicine. 1<sup>st</sup> ed. Gloucester: British Small Animal Veterinary Association. pp. 59-79.

627 Zucca, .D, Redaelli, V., Marelli, S.P., Bonazza, V., Heinzl, E., Verga, M., Luzi, F., 2012. Effect of  
628 handling in pre-weaning rabbits. World Rabbit Sci. 20(2), 97–101.

629

630